

Fig. 4.

*Buliminus stokesi.*

5. BULIMINUS (ENA) CARPATHIUS, n. sp.

Char. Differt a præcedente t. fere perforato-rimata, oblongo-turrita, corneo-lutea, hic illic fuscule strigata, spira convexo-turrita, anfr. 7 distincte convexioribus, celerius accrescentibus,

Fig. 5.

*Buliminus carpathius.*

sutura magis impressa disjunctis, ruguloso-striatis, striis multo distinctioribus, apert. magis obliqua, late acuminato-ovata, perist. simplici, acuto, non labiato, margine columellari prope rimam reflexo, perforationem semitegente.

Alt. 14, diam. min. $4\frac{1}{2}$, maj. $5\frac{1}{2}$ mm.; alt. apert. 5, lat. apert. $3\frac{1}{2}$ mm.

Hab. Karpatho insulam.

Obs. Affinis etiam *Bul. (Enæ) subtili*, Rssm., Cattaroensi, sed t. magis conico-turrita nec fusiformi-turrita et apert. nullomodo labiata.

5. On a new Land-Shell from the New Hebrides.

By JOHN H. THOMSON, C.M.Z.S.

[Received December 5, 1884.]

A shell, of which I can find no published description, was sent me in a package of rare shells collected on the island of Vate, New Hebrides, by Edgar Leopold Layard, Esq., F.Z.S., H.B.M. Consul at Noumea, New Caledonia. Its appearance is at first strikingly similar to that of the North-American *Hyalina arborea*, Say, except as regards

